



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/00, 9/20, 9/70	A1	(11) International Publication Number: WO 98/09613 (43) International Publication Date: 12 March 1998 (12.03.98)
(21) International Application Number: PCT/EP97/04095 (22) International Filing Date: 28 July 1997 (28.07.97) (30) Priority Data: 60/025,444 4 September 1996 (04.09.96) US 08/897,942 21 July 1997 (21.07.97) US (71)(72) Applicant and Inventor: DEGHENGHI, Romano [IT/CH]; Cheseaux Dessus B1, CH-1264 Saint Cergue (CH). (74) Agent: SPADARO, Marco; Bianchetti Bracco Minoja S.r.l., Via Rossini, 8, I-20122 Milano (IT).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PROCESS TO MANUFACTURE IMPLANTS CONTAINING BIOACTIVE PEPTIDES (57) Abstract <p>A process for manufacturing a pharmaceutical composition for the delivery of an effective amount of a bioactive peptide or peptide analog over a period of 1 to 12 months. This process includes the steps of grinding a copolymer of lactic acid and glycolic acid having a ratio of glycolide to lactide units of from about 0 to 5:1 to a particle size of between about 50 and 150 μm; sterilizing the ground copolymer with a dose of between about 1 and 2.5 Mrads of ionizing γ-radiation; wetting the ground and sterilized copolymer with a sterile aqueous slurry of a bioactive peptide or peptide analog; aseptically blending the copolymer and the slurry to obtain a homogeneous mixture of the copolymer and between about 10 and 50 % of the bioactive peptide or peptide analog; drying the mixture at reduced pressure and at temperature not exceeding 25 °C; aseptically extruding the dried mixture at a temperature between about 70 and 110 °C; and aseptically cutting cylindrical rods of about 1 to 2 mm diameter and between about 10 and 25 mm in length from the extruded mixture to form the pharmaceutical implants.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

PROCESS TO MANUFACTURE IMPLANTS CONTAINING BIOACTIVE PEPTIDESTechnical Field

The invention relates to a novel process for preparing
5 implants of bioactive peptides or peptide analogs where such
implants have a more uniform distribution of peptide or
peptide analog therein.

Background Art

10 A wide variety of bioactive peptides and peptide analogs
have been used as active agents for the treatment of various
conditions. These active agents are generally administered in
connection with a polymeric delivery system to control the
release of the agent. For example, peptide analogs of the
15 natural hypothalamic hormone LHRH (Luteinizing Hormone
Releasing Hormone, a decapeptide) are of therapeutic value
when administered for a prolonged period of time with the
appropriate delivery system. Commercially successful delivery
systems include microspheres, microcapsules, microgranules and
20 other implant forms which, when injected subcutaneously or
intramuscularly, release the LHRH analog from a biocompatible
and biodegradable matrix. The matrix is frequently a
copolymer of lactic and glycolic acid ("PLGA", polylactic
glycolic acid) as described, for example, in U.S. Patents
25 3,773,919, 3,887,499, 4,675,189, 4,767,628 and many others.

It has been assumed that a continuous or monophasic
release of such bioactive agents is a highly desirable feature
of such formulations (see, e.g., U.S. Patent 5,366,734). In
fact, it has now been realized that what is really needed is
30 to have the "therapeutic" effect of the peptide or peptide
analog be maintained or sustained over a relatively long span
of time (e.g., three to six months or longer). Thus,
improvements in this area are desired and necessary.

35 Summary of the Invention

The present invention relates to a process for
manufacturing pharmaceutical implants for the delivery of an

effective amount of a bioactive peptide or peptide analog over a period of 1 to 12 months which comprises: grinding a copolymer of lactic acid and glycolic acid having a ratio of glycolide to lactide units of from about 0 to 5:1 to a
5 particle size of between about 50 and 150 μ m; wetting the ground and copolymer with an aqueous slurry of a bioactive peptide or peptide analog; blending the copolymer and the slurry to obtain a homogeneous mixture of the copolymer and between about 10 and 50% of the bioactive peptide; drying the
10 mixture at reduced pressure and at a temperature not exceeding 25°C; extruding the dried mixture at a temperature between about 70 and 110°C; and cutting cylindrical rods of about 1 to 2mm diameter and between about 10 and 25mm in length from the extruded mixture to form the implants.

15 Advantageously, the ground copolymer is sterilized with a dose of between about 1 and 2.5 Mrads of ionizing γ -radiation before being combined with the bioactive peptide, and the blending, extruding and cutting steps are conducted under aseptic conditions. Also, the implants are generally
20 sterilized in a conventional manner prior to being administered to the subject or patient.

The polymers or copolymers form a biodegradable matrix within which is contained a uniform distribution of the peptide or peptide analog. In these copolymers, an
25 advantageous ratio of glycolide to lactide units ranges from about 0.5:1 to 3:1. One particularly preferred copolymer to be used is soluble in benzene and has an inherent viscosity of from 0.51 to 1 (1% in benzene). The amount of slurry is preferably controlled so that the amount of water in the
30 mixture is between about 35 and 65 ml. per 100 grams copolymer, so that the amount of bioactive peptide in these rods is between about 10 to 50 percent by weight.

The bioactive peptide or peptide analog may be an agonist or antagonist of LHRH, GnRH, growth hormone releasing hormone,
35 growth hormone releasing peptide, angiotensin, bombesin, bradykin, cholecystokinin, enkephalin, neurokinin, tachykinin or substance P. The bioactive peptide may also be an

inhibitor such as a renin inhibitor, a protease inhibitor, a metallopeptidase inhibitor, enkephalinase and atrial or brain natriuretic factor degrading enzyme inhibitor. The LHRH analog is preferably a pharmaceutically acceptable salt of an
5 LHRH agonist or antagonist, such as a pharmaceutically acceptable salt of leuprolide, goserelin, triptorelin, buserelin, avorelin, deslorelin, histrelin, cetrorelix, teverelix, ramorelix, antide, nictide, azaline B, azaline C or ganirelix.

10 Another aspect of the invention relates to the pharmaceutical implants obtained according to the process defined herein. These implants are preferably contained in an implanter device with a retractable needle so that they are suitable for subcutaneous injection under the skin of a
15 mammal.

Brief Description of the Drawings

Figure 1 is a graph of serum testosterone and plasma avorelin levels of male beagle dogs for up to 180 days after
20 injection of the avorelin implants of Example 1 of the invention; and

Figures 2 and 3 are graphs of serum LH, FSH and testosterone levels in male patients for up to 33 to 35 weeks after injection of the avorelin implants of Examples 2 and 3
25 of the invention.

Detailed Description of the Preferred Embodiments

Any polylactide polymer or PLGA copolymer can be used to form the biodegradable matrix of this invention. These
30 materials are well known to one of ordinary skill in the art, e.g., in the U.S. patents mentioned above, and need not be further discussed herein. The particular copolymer is selected and then is ground to a particle size of between about 50 and 150 μ m. This grinding step is also conventional
35 and needs no further explanation.

In the most preferred method, the ground copolymer is sterilized with a dose of between about 1 and 2.5 Mrads of

ionizing γ -radiation, again in a conventional manner that is well known to one of ordinary skill in the art.

The ground and sterilized copolymer particles are then wetted with a sterile aqueous slurry of an active agent of a bioactive peptide or peptide analog. This slurry is made by combining the peptide, analog, or a pharmaceutically acceptable salt thereof in sterile water. The amount of the active agent can vary over a wide range, from e.g. about 5 to 50 and preferably about 10 to 25 grams per liter. The solution is then sterilized in a conventional manner, such as by passage through a sterilizing filter. If necessary, the solution can be concentrated to increase the amount of peptide or peptide analog therein. The concentration of the peptide or peptide analog in the solution can be varied to change the resulting dosage of the implant.

Next, the copolymer and the slurry are aseptically blended to obtain a homogeneous mixture of the copolymer and the active agent. Depending upon the desired formulation, the active agent represents between about 10 and 50% and preferably about 15 to 25% of the mixture. As noted above, a water content of about 35 and 65 ml. and preferably about 45 to 55 ml. per 100 grams copolymer in the mixture is desired. Next, the mixture is dried at reduced pressure and at a temperature not exceeding 25°C to form the pharmaceutical composition. If necessary, this composition can be formulated with conventional carriers as a suspension for injection.

Alternatively, the dried composition can be extruded with a conventional extrusion device at a temperature between about 70 and 110°C to form a "spaghetti" or continuous rod product. The use of heat in the extrusion step helps further dry the product. To form the implants, these cylindrical rods are aseptically cut into pieces of about 1 to 2mm diameter and between about 10 and 25mm in length from the extruded mixture. The length of the implant is another mechanism for varying the dosage of bioactive peptide or peptide analog therein. These products can then be implanted subcutaneously beneath the skin of the patient using conventional implanting devices.

The present invention provides an effective release (i.e. in terms of therapeutic effectiveness) of a bioactive peptide, or peptide analog, such as an LHRH analog, even if such release, as measured by plasma level of the peptide, or
5 peptide analog, is intermittent or discontinuous. This effectiveness can be achieved, for example, by the internalization or down-regulation of pituitary receptors following their exposure to LHRH agonists or to LHRH antagonists which are intrinsically long acting.

10 The process of this invention can be applied to a wide variety of peptides or peptide analogs. In addition to LHRH analogs herein mentioned, GnRH or growth hormone releasing hormones or peptides can be mentioned. Generally, any peptides or peptide analogs that are chemically stable under
15 the process conditions and that provide a sustained delivery is desirable from a therapeutic point of view. Non-limiting examples of such peptides and peptide analogs are somatostatin and somatostatin analogs, agonist and antagonist analogs of angiotensin II, bombesin analogs, preferably bombesin
20 antagonists, bradykinin antagonists, preferably with minimal histamine releasing properties, cholecystokinin analogs, preferably cholecystokinin antagonists, enkephalin analogs, neurokinins, tachykinins and substance P antagonists, renin inhibitors and other aspartyl protease inhibitors, such as HIV
25 protease inhibitors, metallopeptidase inhibitors, such as angiotensin converting enzyme, enkephalinase and atrial or brain natriuretic factor degrading enzyme inhibitors. The skilled artisan will favor those peptides and peptidomimetic compounds which are not, or are poorly, absorbed by the oral
30 route in animals and humans, and will adjust the dose of the compound to be formulated in the implants of the present invention according to the biological potency of such compound, the necessary daily effective dose and the estimated duration of release from the formulation.

35 The present invention also eliminates contamination of such formulations with organic solvents, particularly chlorinated ones, such as chloroform or methylene chloride,

which are typically utilized in the manufacture of microspheres or microcapsules by the coacervation-solvent evaporation methods (see, e.g., U.S. Patent 3,773,919) or which are used to sterilize PLGA copolymers by filtration.

5 The present invention does not make use of any organic solvent, but takes advantage of the unorthodox use of water, a solvent hitherto considered unsuitable for such formulations because of its deleterious effect on the polyester (copolymer) of the PLGA matrix, where it can accelerate chemical
10 hydrolysis and also damage structural integrity upon exposure to ionizing radiation (formation of free radicals) during the sterilization step necessary for safety considerations.

Another advantage of this unorthodox use of water is to achieve a uniform coating of the active principle on the
15 granulated polymer powder, resulting in a much needed and highly desired uniformity of the mixture, an essential condition of the manufacturing process. A further unexpected advantage of this unconventional solvent is the "wettability" of the powdery mixture which would otherwise create serious
20 problems due to formation of static electrical charges which can cause unacceptable mechanical losses and loss of uniformity.

The instant process further provides with a simple method of sterilization of the composition by subjecting the polymer
25 to ionizing radiation prior to blending the polymer with the bioactive peptides or peptide analogs which are invariably damaged by such radiation, resulting in unwanted byproducts. A further advantage of the instant process is to provide a variable sterilizing dose of radiation (from 1 to 2.5 Mrad)
30 predetermined by the actual biomass present in the co-polymer, with a resulting safety without undue creation of radiolysis artifacts.

Examples

35 The following examples are submitted to illustrate the effectiveness of the most preferred formulations of the invention.

Example 1

The manufacturing process is conducted in a commercially available isolator (ARFL, Neuilly-sur-Marne, France) equipped with air-locks for the introduction of pre-sterilized
5 components and itself sterilized by previous peracetic acid treatment. The extrusion machine is a commercially available single screw extruder (Brabender, 47055 Duisburg, Germany) equipped with pressure and temperature probes. The cutting machine is commercially available (Davis-Standard Corp. Cedar
10 Grove, N.J., USA). Blenders/mixers and weighing instruments are conventional equipment.

A quantity of 80g of racemic lactic acid and glycolic acid copolymer (75:25) soluble in benzene and of inherent viscosity of 0.60 (1% in benzene) (PuracBiochem B.V.,
15 Gorinchem, Netherlands) is ground and sieved to collect the fraction of particles between 50 to 150 μm and sterilized with an ionizing γ -radiation of 1.5 Mrads by a commercial laboratory (Caric-Mediris, Fleurus, Belgium) and introduced through the air-lock into the sterile isolator.

20 Separately, 23 grams of the LHRH analog avorelin acetate (INN), or (2Methyl-D-Trp)⁶(des-Gly)¹⁰(ProEthylamide)⁹LHRH acetate, dissolved in 500ml of sterile water and filtered through a Millipore 0.2 μm sterilizing filter. The sterile solution is reduced by evaporation to a volume of 50ml and the
25 resulting mixture is dispersed through the ground co-polymer. The wet mixture is blended to obtain a granulate containing 20% of avorelin. Such mixture is dried at 25°C under reduced pressure and then extruded at a temperature gradient from 70 to 110°C at pressures of 3500 p.s.i. The resulting extrudate
30 is aseptically cut to give rods of 1.5mm diameter and 15mm long, containing 10 mg avorelin, which are inserted into a pre-sterilized implanter with a retractable needle (SFM GmbH, D-6480 Wächtersbach, Germany) sealed and used as such, or optionally further sterilized with a dose of 1.5 Mrad of γ -
35 radiation before clinical use.

When implanted s.c. into male beagle dogs, after the initial stimulation of LH and testosterone, castration levels

of testosterone were maintained for 6 months. The plasma levels of avorelin, after a short-lived burst, fell to a nadir at 40 days and rose again at 120 days before becoming undetectable at day 160. These results are shown in Figure 1.

5

Example 2

Following essentially the procedure of Example 1, 10 mg avorelin implants were prepared, further sterilized, and implanted into healthy male patients. After initial
10 stimulation of LH, FSH and testosterone, these levels were significantly reduced, with the testosterone level being maintained below a castration level for 33 weeks. These results are shown in Figure 2.

15 Example 3

Following essentially the procedure of Example 2, except that the length of the implant was increased to provide a dose of avorelin of 15 mg was prepared. These implants were sterilized and implanted into healthy male patients. After
20 initial stimulation of LH, FSH and testosterone, these levels were significantly reduced, with the testosterone level being maintained below a castration level for 33 weeks. These results are shown in Figure 3.

25 Example 4

Following essentially the procedure of Example 1 with appropriate modifications required by the individual LHRH analog, rods containing 22mg of leuprolide, 10mg of goserelin and 30mg of teverelix were similarly obtained.

30

35

Claims

1. A process for manufacturing pharmaceutical implants
5 for the delivery of an effective amount of a bioactive peptide
or peptide analog over a period of 1 to 12 months which
comprises:

grinding a copolymer of lactic acid and glycolic acid
having a ratio of glycolide to lactide units of from about 0
10 to 5:1 to a particle size of between about 50 and 150 μ m;

wetting said ground and sterilized copolymer with a
sterile aqueous slurry of a bioactive peptide or peptide
analog;

blending the copolymer and the slurry to obtain a
15 homogeneous mixture of said copolymer and between about 10 and
50% of the bioactive peptide or peptide analog;

drying said mixture at reduced pressure and at
temperature not exceeding 25°C;

extruding said dried mixture at a temperature between
20 about 70 and 110°C; and

cutting cylindrical rods of about 1 to 2mm diameter and
between about 10 and 25mm in length from the extruded mixture
to form the pharmaceutical implants.

25 2. The process of claim 1 which further comprises
sterilizing said ground copolymer with a dose of between about
1 and 2.5 Mrads of ionizing γ -radiation before adding the
aqueous slurry thereto.

30 3. The process of claim 1 which further comprises
conducting the blending, extruding and cutting steps are
conducted aseptically.

4. The process of claim 1 which further comprises
35 selecting the copolymer to be used to be one which is soluble
in benzene and has an inherent viscosity of from 0.51 to 1 (1%
in benzene).

5. The process of claim 1 wherein the amount of slurry is controlled so that the amount of water in the mixture is between about 35 and 65 ml. per 100 grams copolymer.

5 6. The process of claim 1 wherein the amount of slurry is controlled so that the amount of bioactive peptide or peptide analog in the rods is between about 10 to 50 percent by weight.

10 7. The process of claim 1 wherein the ratio of glycolide to lactide units in the copolymer ranges from about 0.5:1 to 3:1

8. The process of claim 1 wherein the bioactive peptide
15 or peptide analog is an agonist or antagonist of LHRH, GnRH, growth hormone releasing hormone, growth hormone releasing peptide, angiotensin, bombesin, bradykin, cholecystokinin, enkephalin, neurokinin, tachykinin or substance P.

20 9. The process of claim 1 wherein the bioactive peptide or peptide analog is a renin inhibitor, a protease inhibitor, a metallopeptidase inhibitor, enkephalinase and atrial or brain natriuretic factor degrading enzyme inhibitor.

25 10. The process of claim 8 wherein the bioactive peptide or peptide analog is a pharmaceutically acceptable salt of leuprolide, goserelin, triptorelin, buserelin, avorelin, deslorelin, histrelin, cetrorelix, teverelix, ramorelix, antide, nictide, azaline B, azaline C or ganirelix.

30

11. A pharmaceutical implant obtained according to the process of any one of the preceding claims.

12. The pharmaceutical implant of claim 11 contained in
35 an implanter device with a retractable needle and suitable for subcutaneous injection under the skin of a mammal.

1/3




FIG. 1




FIG. 2




FIG. 3

INTERNATIONAL SEARCH REPORT

International Application No

PC1/EP 97/04095

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K9/00 A61K9/20 A61K9/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 38 22 459 A (BPD BIOPHARM DEV LTD) 30 March 1989 ---	
A	GB 2 234 169 A (DEBIOPHARM SA) 30 January 1991 ---	
A	CH 685 285 A (DEBIO RECH PHARMA SA) 31 May 1995 ---	
A	GB 2 249 725 A (DEBIO RECH PHARMA SA) 20 May 1992 ---	
A	US 5 456 917 A (WISE DONALD L ET AL) 10 October 1995 -----	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

20 January 1998

Date of mailing of the international search report

03/02/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Fischer, W

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/EP 97/04095

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 3822459 A	30-03-89	GB 2209937 A	01-06-89
		AT 397035 B	25-01-94
		AU 2232688 A	23-03-89
		BE 1001685 A	06-02-90
		CA 1326438 A	25-01-94
		CH 675968 A	30-11-90
		DK 518988 A	22-03-89
		FI 96919 B	14-06-96
		FR 2620621 A	24-03-89
		GR 88100619 A	22-06-89
		IE 60608 B	27-07-94
		JP 1121222 A	12-05-89
		JP 1981991 C	25-10-95
		JP 7013023 B	15-02-95
		LU 87340 A	06-04-89
		NL 8802323 A	17-04-89
		NO 178604 B	22-01-96
		SE 503406 C	10-06-96
		SE 8803321 A	22-03-89
		US 5192741 A	09-03-93
GB 2234169 A	30-01-91	CH 679207 A	15-01-92
		AT 397197 B	25-02-94
		AU 619996 B	06-02-92
		AU 5910390 A	31-01-91
		BE 1003093 A	19-11-91
		CA 2021767 A,C	29-01-91
		DE 4023134 A	31-01-91
		DK 180790 A	29-01-91
		FI 97688 B	31-10-96
		FR 2650182 A	01-02-91
		GR 1001215 B	21-06-93
		IE 65397 B	18-10-95
		IL 95120 A	07-10-94
		IT 1243357 B	10-06-94
		JP 1933192 C	26-05-95
		JP 3066625 A	22-03-91
		JP 6062427 B	17-08-94
		LU 87772 A	11-12-90
		NL 9001646 A	18-02-91

INTERNATIONAL SEARCH REPORT

Information on patent family members

I national Application No

PCT/EP 97/04095

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2234169 A		NO 300304 B PT 94842 B SE 504279 C SE 9002522 A US 5134122 A US 5439688 A US 5225205 A	12-05-97 30-04-97 23-12-96 29-01-91 28-07-92 08-08-95 06-07-93
CH 685285 A	31-05-95	NONE	
GB 2249725 A	20-05-92	CH 681425 A AT 223591 A,B BE 1004923 A CA 2055115 A DE 4136930 A ES 2049617 A FR 2668707 A IT 1252870 B NL 9101877 A SE 506448 C SE 9103348 A US 5439688 A	31-03-93 15-07-93 23-02-93 15-05-92 20-08-92 16-04-94 07-05-92 28-06-95 01-06-92 15-12-97 15-05-92 08-08-95
US 5456917 A	10-10-95	EP 0693923 A JP 8512288 T WO 9423698 A	31-01-96 24-12-96 27-10-94